Salinity Science Results from the Gulf of Mexico (GOM)


Jorge Vazquez


NASA Jet Propulsion Laboratory

Scientist for Sea Surface Temperature and Salinity

Plotting SMAP data in LAS

- Following slides show how to plot SMAP data using one particular tool available from PO.DAAC: Live Access Server (LAS)
- LAS can be used to compare SMAP data:
 - RSS 40 km 8-day running mean
 - RSS 70 km 8-day running mean
- It is also capable of generating time series of averaged data (e.g., over an area)





You will prompted to select a dataset

US Department of Commerce NOAA | OAR | PMEL | Contacts | Privacy Policy | Disclaimer | mailto:


Compute: None

My selections:

Select:


Live Access Server

PO.DAAC LAS V8.6.1



Live Access Server PO.DAAC LAS V8.6.1 Print... Link... Animate Correlation Viewer Google Earth Show Values Export to Desktop Application Save As... Update Plot One Plot Annotations X DATASET: RSS SMAP Level 3 Sea Surface Salinity Standard Mapped Image 8-Day Running Mean V2.0 Validated Dataset @ VARIABLE: SMAP sea surface salinity (1e-3) Plot Options TIME: 31-MAR-2015 12:00 OPeNDAP URL: http://thredds.jpl.nasa.gov/thredds/dodsC/ncml_aggregation/SalinityDensity/smap/aggregate__SMAP_RSS_L3_SSS_SMI_8DAY-RUNNINGMEAN_V2.ncml . Subsampled 4 in X, 4 in Y LAS 8./Ferret 6.96 NOAA/PMEL Date/Time: 2015 Mar 31 3 SMAP sea surface salinity 90 N 31.5 32.5 33.1 33.3 33.5 33.7 33.9 34.1 34.3 34.5 34.7 34.9 35.1 35.3 35.5 35.7 35.9 36.5 37.5 1.192 90 S CE Compute: Maps Latitude-Longitude Line Plots Time Longitude Latitude Hovmoller Plots Longitude-time Latitude-time 80°S -By default, you will see a


global data map US Department of Commerce NOAA | OAR | PMEL | Contacts | Privacy Policy | Disclaimer | mailto:




Pick a specific area by (1) entering coordinates or (2) drawing a box.

Click Update Plot (3) to see your results.


US Department of




40 km SMAP data for 31-Mar-15



Change the date using pull-down menus Click Update Plot to see your results.



40 km SMAP data for 31-Jul-16




Note: You can click "Plot Options" and choose "Off" to view non-interpolated data

US Depa

GOM – 40 km RSS Data

(interpolated)


Data are close to the coast but noisy




GOM - 70 km RSS Data

(interpolated)

70-km
data are
smoother
but further
from the
coast






To generate time-series average data, use the "Compute" pull-down menu...

Live Access Server

...and set "Start date/time" & "End date/time." Click Update Plot to see your results.



Average GOM salinity Jan – Dec 2016.


(1) You can also compare with other data using the "Compare 2" option.

16

Time Series – GOM Average SMAP 40 km data set 36.20 36,00 35,80 35.60 JUL DEC AUG OCT NOA 2016 36,40 SMAP 70 km data set 36.20 35,80 35.60 DEC JAN AUG NOV 2016

SMAP 70 km Data – Amazon

• SMAP data have also been used to create a timeseries visualization of the Amazon Plume



Recap of LAS Instructions

- http://podaac.jpl.nasa.gov => Go to to drop-down menu under "Data Access". Click on "LAS (L3 subsetting)"
- Go to "Data Set" (RSS 40-km & 70-km and JPL 60-km SMAP Salinity data sets available at 8-day running mean and monthly resolutions)
- Click on the desired data set
- To subset, put in coordinates to define desired box
 - Underneath the map on the left hand side of LAS interface
- Select time range
- Select desired output (lat-lon map; time series; Hovmoller diagrams)
- At the top right corner, clicking on "Save As..." gives you option to save output as: NetCDF; ASCII; CSV; arcGrid.
- Other tools include Animate, Correlation Viewer, etc.



https://salinity.oceansciences.org

https://podaac.jpl.nasa.gov/SeaSurfaceSalinity

http://www.remss.com/missions/smap/salinity/